

The usual mammalian dosage range for a 70kg human subject is from 5 to 500mg per day (.07mg to 7.1mg per kg of weight per day), preferably 15 to 120mg per day (.2mg to 1.7mg per kg of weight per day), optionally in divided portions. Therapeutic agents of this type are generally prescribed for a specific time span, such as for a five day period depending upon the disease state being treated."

Page 9 (bottom), 10 (top), delete last paragraph:

"Thus choriocarcinoma type states may be treated with 15mg to 60mg per day for a five day period and repeated three to five times as required while mycosis fungoides type condition may require lower doses, such as 5mg to 25mg daily over greater periods of time which may be measured in weeks or months."

Page 10, delete first complete paragraph:

"In addition, the compounds of this invention may be employed in combination or conjunction with other agents, such as cyclophosphamide, vincristine, cytarabine, fluorouracil, mercaptopurine, prednisone, lomustine, etc. When used in combination or conjunction with the foregoing agents, the dosage may be lowered."

In the Claims:

Cancel claims 4-9 without prejudice to their reentry at some later date.

Please add the following new claims:

10. ~~The compound of claim 1 being in crystalline form.~~
11. The compound of claim 2 being in crystalline form.
12. ~~The compound of claim 1 being in powdery form.~~
13. The compound of claim 3 being in powdery form.